



## Marine Vibrionaceae as a source of bioactive natural products

Månsson, Maria; Wietz, Matthias; Gram, Lone; Larsen, Thomas Ostfeld

*Publication date:*  
2011

*Document Version*  
Early version, also known as pre-print

[Link back to DTU Orbit](#)

*Citation (APA):*  
Månsson, M., Wietz, M., Gram, L., & Larsen, T. O. (2011). *Marine Vibrionaceae as a source of bioactive natural products*. Abstract from 7th European Conference on Marine Natural Products, Strömstad, Sweden.  
<http://www.fkog.uu.se/7ecmnp/>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Marine *Vibrionaceae* as a source of bioactive natural products

Maria Månsson<sup>1</sup>, Matthias Wietz<sup>2</sup>, Lone Gram<sup>3</sup>, Thomas Ostenfeld Larsen<sup>4</sup>

<sup>1</sup>maj@bio.dtu.dk; <sup>2</sup>mwie@food.dtu.dk; <sup>3</sup>gram@food.dtu.dk; <sup>4</sup>tol@bio.dtu.dk

*Vibrionaceae* are Gram-negative bacteria found widespread in the marine environment where they are particularly abundant on the surface of marine macroorganisms. Production of antibacterial compounds appears to be common among vibrios, yet vibrios are largely underexplored for their proclivity to produce secondary metabolites.

We have studied the production of antibacterial compounds in *Vibrionaceae* collected during a global marine expedition, Galathea 3. Apart from growth inhibitory compounds, we searched for compounds that interfere with virulence regulation in *Staphylococcus aureus*.

We found that some strains were capable of producing antibacterial compounds when grown on natural substrates such as chitin or seaweed. One *Vibrio coralliilyticus* strain was capable of producing the antibacterial compound when using chitin as the sole carbon source and in a live chitin model system, suggesting an ecological function.

Using chemical profiling, vibrio strains were compared on a global scale, revealing that the production of certain compounds is a conserved feature independent of sample locations. Chemical screening techniques such as explorative solid-phase extraction led to the isolation of two novel depsipeptides, solonamide A and B, as potent inhibitors of the *agr* QS system involved in virulence expression in *S. aureus*. Of special interest was a pronounced effect against a highly virulent, CA-MRSA strain (USA300).

In conclusion, we found that vibrios are competent producers of secondary metabolites, some of which possess biological activities attractive for alternative strategies in antibacterial therapy.